

Cell Lines Available

FIBROBLAST NON-CLASSIC MUTATIONS					
PSADFN485 ⁵	Proband	4 yrs 5 mos	Male	LMNA heterozygous c.412G>A; (p.E138K)	
PSADFN425 ^{1, 5}	Proband	20 yrs 11 mos	Male	LMNA Exon 1, heterozygous c.331G>A (p.Glu111Lys) Intron 6,1158-44 C>T	PSALBV295
PSADFN412 ^{1, 2, 5}	Proband	7 yrs 1 mo	Male	LMNA Exon 11, heterozygous c.1762T>C (p.C588R)	
PSADFN328 ^{1, 2, 5} (Cells grow poorly)	Proband	12 yrs 5 mos	Female	LMNA Exon 11, heterozygous c.1822 G->A (p.Gly608Ser)	PSALBV296
PSADFN257 ^{2, 5}	Proband	1 yr 10 mos	Male	LMNA Exon 10, homozygous c.1619 T>C (p.Met540Thr)	
PSADFN086 ^{2, 5} (Cells grow poorly)	Proband	0 yrs 7 mos	Male	LMNA Exon 11, c.1968+1 G>A	PSALBV083
PSADFN542	Proband	75 yrs 2 mos	Male	LMNA heterozygous Exon 11, c.1930 C>T (p.Arg644Cys)	
PSADFN576	Proband	34 yrs 11 mos	Female	LMNA Exon 11/Intron 11 junction, heterozygous c. 1968+5G>A	

¹Representative cultures from this cell line have tested negative for mycoplasma contamination at The PRF Cell and Tissue Bank. Please note: mycoplasma testing is performed on random passages. As part of our on-going quality control, The PRF Cell and Tissue Bank periodically tests the cultures for mycoplasma contamination using R&D Systems Mycoplasma Detection Kit (catalog # CUL001B).

²Histograms of mutational analysis sequenced by the PRF Cell and Tissue Bank available.

³Representative cultures from this cell line have tested negative for mycoplasma contamination at Rutgers University Cell and DNA Repository via real time PCR assay.

⁴Representative cultures from this cell line have tested negative for mycoplasma contamination at Ottawa Hospital Research Institute. Please note: mycoplasma testing is performed on random passages. As part of our on-going quality control, The Human Pluripotent Stem Cell Facility/Dr. William Stanford laboratory periodically tests the cultures for mycoplasma contamination using a PCR based approach (Detection of mycoplasma contaminations., Uphoff CC, Drexler HG., Methods Mol Biol. 2013;946:1-13. doi: 10.1007/978-1-62703-128-8_1. PMID:23179822).



FAMILY NON-CLASSIC FIBROBLAST CELL LINES					
Cell Line #	Relation to Proband	Age at Donation	Gender	Mutational Analysis ²	Other Lines From This Donor
PSADFN489	Proband (mild progeroid phenotype)	3 yrs 2 mos	Female	LMNA Exon 11, heterozygous c.1824C>T (p.Gly608Gly); and SMC3 c.562A>G, p.K188E	
PSFDFN714	Father of HGADFN489	38 yrs 6 mos	Male	Negative	
PSMDFN713	Mother of HGADFN489	35 yrs 11 mos	Female	Negative	
PSADFN392	Proband	7 yrs 4 mos	Male	LMNA Exon 11, c.1968+2T>C	
PSFDFN394 ¹	Father of PSADFN392	49 yrs 1 mo	Male	Not performed	
PSMDFN393 ¹	Mother of PSADFN392	44 yrs 8 mos	Female	Not performed	
PSADFN373 ¹	Proband	5 yrs 9 mos	Male	ZMPste24 Exon 10, homozygous c.1274T>C (p.Leu425Pro)	PSALBV341
PSFDFN376 ¹	Father of PSADFN373	32 yrs 6 mos	Male	ZMPste24 Exon 10, heterozygous c.1274T>C (p.Leu425Pro)	PSFLBV344
PSMDFN375 ¹	Mother of PSADFN373	32 yrs 9 mos	Female	ZMPste24 Exon 10, heterozygous c.1274T>C (p.Leu425Pro)	PSMLBV343
PSADFN363 ^{1, 2, 5}	Proband	8 mos	Male	LMNA Exon 6, heterozygous c.973G>A (p.Asp325Asn)	
PSFDFN365 ^{1, 2, 5}	Father of PSADFN363	44 yrs 2 mos	Male	Not performed	
PSMDFN364 ^{1, 2, 5}	Mother of PSADFN363	36 yrs 10 mos	Female	Not performed	

Cell Lines Available

¹Representative cultures from this cell line have tested negative for mycoplasma contamination at The PRF Cell and Tissue Bank. Please note: mycoplasma testing is performed on random passages. As part of our on-going quality control, The PRF Cell and Tissue Bank periodically tests the cultures for mycoplasma contamination using R&D Systems Mycoplasma Detection Kit (catalog # CUL001B).

²Histograms of mutational analysis sequenced by the PRF Cell and Tissue Bank available.

³Representative cultures from this cell line have tested negative for mycoplasma contamination at Rutgers University Cell and DNA Repository via real time PCR assay.

⁴Representative cultures from this cell line have tested negative for mycoplasma contamination at Ottawa Hospital Research Institute. Please note: mycoplasma testing is performed on random passages. As part of our on-going quality control, The Human Pluripotent Stem Cell Facility/Dr. William Stanford laboratory periodically tests the cultures for mycoplasma contamination using a PCR based approach (Detection of mycoplasma contaminations., Uphoff CC, Drexler HG., Methods Mol Biol. 2013;946:1-13. doi: 10.1007/978-1-62703-128-8_1. PMID:23179822).



Cell Lines Available

FAMILY NON-CLASSIC FIBROBLAST CELL LINES CONTINUED					
Cell Line #	Relation to Proband	Age at Donation	Gender	Mutational Analysis	Other Lines From This Donor
PSMDFN346 ^{2, 5}	Mother of PSADFN345 (proband line not available)	21 yrs 10 mos	Female	LMNA Exon 11, Negative	
PSADFN325 ^{1, 2, 5}	Proband	6 yrs 9 mos	Male	LMNA Exon 11/Intron 11 junction, heterozygous c. 1968+5G>C	
PSFDFN327 ^{2, 5}	Father of HGADFN325	36 yrs 3 mos	Male	Not performed	
PSMDFN326 ^{2, 5}	Mother of HGADFN325	36 yrs 10 mos	Female	Not performed	
PSADFN317 ⁵	Proband (& sibling of PSADFN318)	3 yr 9 mo	Male	ZMPste24 Exon 6, heterozygous c.743C>T(p.Pro248Leu); Exon 10, heterozygous c.1349G>A(p.Trp450Stop)	
PSADFN318 ^{1, 5}	Proband (& sibling of PSADFN317)	5 mos	Male	ZMPste24 Exon 6, heterozygous c.743C>T(p.Pro248Leu); Exon 10, heterozygous c.1349G>A(p.Trp450Stop)	
PSFDFN319	Father of PSADFN317 & PSADFN318	39 yrs 0 mo	Male	Not performed	
PSMDFN320 ¹	Mother of PSADFN317 & PSADFN318	36 yrs 8 mo	Female	Not performed	

¹Representative cultures from this cell line have tested negative for mycoplasma contamination at The PRF Cell and Tissue Bank. Please note: mycoplasma testing is performed on random passages. As part of our on-going quality control, The PRF Cell and Tissue Bank periodically tests the cultures for mycoplasma contamination using R&D Systems Mycoplasma Detection Kit (catalog # CUL001B).

²Histograms of mutational analysis sequenced by the PRF Cell and Tissue Bank available.

³Representative cultures from this cell line have tested negative for mycoplasma contamination at Rutgers University Cell and DNA Repository via real time PCR assay.

⁴Representative cultures from this cell line have tested negative for mycoplasma contamination at Ottawa Hospital Research Institute. Please note: mycoplasma testing is performed on random passages. As part of our on-going quality control, The Human Pluripotent Stem Cell Facility/Dr. William Stanford laboratory periodically tests the cultures for mycoplasma contamination using a PCR based approach (Detection of mycoplasma contaminations., Uphoff CC, Drexler HG., Methods Mol Biol. 2013;946:1-13. doi: 10.1007/978-1-62703-128-8_1. PMID:23179822).



Cell Lines Available

FAMILY NON-CLASSIC FIBROBLAST CELL LINES CONTINUED						
Cell Line #	Relation to Proband	Age at Donation	Gender	Mutational Analysis	Other Lines From This Donor	
PSADFN386 ¹	Proband	11 mos	Female	Mosaic: DNA from Fibroblasts: c.1968 +2T>C DNA from Blood: 4.7% c.1968+2T>C mutation, and 41.3% c.1968+2T>A mutation See reference: Bar DZ, Arlt MF, Brazier JF, et al. A novel somatic mutation achieves partial rescue in a child with Hutchinson-Gilford progeria syndrome. Journal of Medical Genetics 2017;54:212-216		
PSMDFN387 ¹	Mother of PSADFN386	36 yrs 5 mos	Female	LMNA Exon 11, Negative		
PSFDFN388 ¹	Father of PSADFN386	38 yrs 1 mo	Male	LMNA Exon 11, Negative		

¹Representative cultures from this cell line have tested negative for mycoplasma contamination at The PRF Cell and Tissue Bank. Please note: mycoplasma testing is performed on random passages. As part of our on-going quality control, The PRF Cell and Tissue Bank periodically tests the cultures for mycoplasma contamination using R&D Systems Mycoplasma Detection Kit (catalog # CUL001B).

²Histograms of mutational analysis sequenced by the PRF Cell and Tissue Bank available.

³Representative cultures from this cell line have tested negative for mycoplasma contamination at Rutgers University Cell and DNA Repository via real time PCR assay.

⁴Representative cultures from this cell line have tested negative for mycoplasma contamination at Ottawa Hospital Research Institute. Please note: mycoplasma testing is performed on random passages. As part of our on-going quality control, The Human Pluripotent Stem Cell Facility/Dr. William Stanford laboratory periodically tests the cultures for mycoplasma contamination using a PCR based approach (Detection of mycoplasma contaminations., Uphoff CC, Drexler HG., Methods Mol Biol. 2013;946:1-13. doi: 10.1007/978-1-62703-128-8_1. PMID:23179822).